

Contact information

If you need to contact us, please use our contact numbers below:

Monday to Saturday 8am-8pm: Same Day Emergency Care 0131 537 2137

Monday to Saturday 8pm-8am: Medical Assessment Unit 0131 537 1330

Sunday: Medical Assessment Unit 0131 537 1330



Information for patients

Worsening symptom advice for deep vein thrombosis (DVT - a blood clot in a vein)



You have been assessed today in Same Day Emergency Care and your doctor or advanced nurse practitioner has decided that you are safe to go home.

It is important to make sure that you know what to do if you feel less well after leaving the hospital.

If you develop the following symptoms, you should seek medical attention urgently. This will be discussed with you verbally as well.

- Increasing leg swelling or pain, or a change in colour of your leg, particularly if this happens suddenly
- Shortness of breath
- Chest pains
- Coughing up blood
- Feeling faint or dizzy
- Racing heart.

If you develop these symptoms you should phone 999 or go to hospital as soon as possible.

If you have been given blood thinning treatment for returning for an ultrasound scan:

You may have been given a blood thinning injection (dalteparin) or tablets (apixaban) overnight while you await your scan.

- If you develop any unusual or spontaneous bleeding or bruising, you must go to hospital as soon as possible
- If you have a head injury, it is vital that you go to accident and emergency as soon as possible.

Do not undertake any sporting activities or strenuous exercise, have a hot bath/shower, or undergo any massage until after you have had your scan.