

Endoscopy & Oesophageal dilatation

Discharge Information for patients

Upper gastrointestinal endoscopy and oesophageal dilatation with sedation discharge information

Possible side effects you may experience:

Abdominal discomfort: You may feel a little bloated and have some wind-like pains because of the gas passed into your bowel during the procedure. This should settle within a few hours.

Sore throat: You may have a sore throat. This is quite normal and will pass within 24-48hrs. If you had a biopsy taken, you may bring up a small amount of blood-stained fluid.

Complications

Contact NHS 24 or A&E if you experience any of the following symptoms.

- Severe pain in the neck, chest or abdomen
- Bringing up more than a small amount of blood
- Breathing difficulties
- High temperature, sweats or chills
- Passing blood (red or black) in your bowel movements
- Vomiting

Contact telephone numbers can be found overleaf.

Medication

Please continue to take any regular medication unless otherwise advised.

Difficulty in swallowing: After this procedure you will experience some difficulty in swallowing solid food. This should improve within 2-3 days.

Diet

Day of procedure: Follow a sloppy diet e.g. porridge, custard, yogurt etc.

Avoid hot drinks and food (warm/cool is better)

Day 1 after procedure: Your normal diet avoiding abrasive foods such as toast.

Day 2 after procedure: Your normal diet as tolerated

Discomfort at the injection site

If you had a cannula inserted, you may experience slight discomfort at the injection site. If this becomes increasingly red, tender or swollen, please seek advice from your GP

Endoscopy and Oesophageal Dilation V1.0

Approved by: NHS Lothian Patient Information Aug 2023

Review date: Aug 2026

Sedation

Sedation makes you drowsy and can impair your reflexes and judgement. For 24 hours after you have had sedation:

Do not go to work	Do not drink alcohol	
Do not sign any important documents	Do not drive or ride a bike	
Avoid looking after young children	Take care in the bath or shower	
Do not use any kind of machinery, including household appliances		Do not take sleeping tablets

We advise you to rest quietly with adult supervision for the rest of the day.

Results

You should have already received a copy of your results, but a copy of the report will also be sent to your GP. If you had biopsies taken these results will be sent to you as soon as they are available. This can take several weeks. Your GP will also be sent a copy of your biopsy results. If you need to make an appointment with your GP regarding your results, please contact your health centre to ensure they have received them first.

Fol	low	u	p
-----	-----	---	---

Any other relevant information		
If any changes are required following t	his review, you will be inforr	ned at that time.
You may need a follow up procedure. procedure date. This will make sure it guidelines.		
☐ A follow up procedure is required		
□ No further follow up required □	Follow up with your GP	☐ Follow up in clinic

Contact telephone numbers

Western General Hospital	0131 537 1327 or 0131 537 1695
Royal Infirmary of Edinburgh	0131 242 1600
St John's Hospital (voicemail)	01506 523 982
East Lothian Community Hospital	01620 642 885
NHS 24	111 Freephone
Accident and Emergency (Royal Infirmary of Edinburgh switchboard)	0131 536 1000

Endoscopy and Oesophageal Dilation V1.0 Approved by: NHS Lothian Patient Information Aug 2023

Review date: Aug 2026