

Endoscopy with Sedation

Discharge information for patients



Possible side effects of Upper Gastrointestinal Endoscopy:

Abdominal discomfort You may feel a little bloated and have some wind-like pains because of the gas passed into your stomach during the procedure. This should settle within a few hours.

Sore throat You may have a sore throat. This is quite normal and will pass after 24 to 48 hours. If you had a biopsy taken you may bring up a small amount of blood-stained fluid.

Complications

Contact NHS 24 or A&E if you experience any of the following:

- Severe pain in the neck, chest or abdomen
- Bringing up blood, more than a small amount
- Breathing difficulties
- High temperature, sweats or chills
- Passing blood (red or black) in your bowel movements
- Vomiting

Contact numbers can be found on the back of this page.

Medication

Please continue to take any regular medication unless otherwise advised.

Discomfort at injection site

You may experience slight discomfort at the site where the sedative injection was given. If this site becomes increasingly red, tender or swollen, please seek advice from your GP.

Sedation

Sedation makes you drowsy and can impair your reflexes and judgement. For 24 hours after you have had sedation:

Do not go to work	Do not drink alcohol	
Do not sign any important documents		
Avoid looking after young children		
Do not use any kind of machinery, including household appliances		Do not take sleeping tablets

We advise you to rest quietly with adult supervision for the rest of the day.

☐ Follow up with your GP ☐ Follow up in clinic

Results

You should have already received a copy of your results, but a copy of the report will also be sent to your GP.

If you had biopsies taken, these results will be sent to you as soon as they are available. This can take several weeks. Your GP will also be sent a copy of your biopsy results.

If you need to make an appointment with your GP regarding your results, please contact your health centre to ensure they have received them first.

Follow up

☐ A follow up procedure is required	
If you need a follow up procedure as part of a surveillance programme or to reassess any abnorm	alities
found at your endoscopy your case will be reviewed by us 2 months before the procedure date.	Thic

found at your endoscopy, your case will be reviewed by us 2 months before the procedure date. This will make sure it still needs to be done according to the most up-to-date guidelines.

If any changes are required following this review, you will be informed at that time.

Any other relevant information

☐ No further follow up required

Contact telephone numbers

Western General Hospital	0131 537 1327 or 0131 537 1695

Royal Infirmary of Edinburgh 0131 242 1600

St John's Hospital (voicemail) 01506 523 982

East Lothian Community Hospital 01620 642 885

Leith Community Treatment Centre 0131 536 6442

NHS 24 111 Freephone

Accident and Emergency 0131 536 1000

(Royal Infirmary of Edinburgh switchboard)